Mon. Not. R. Astron. Soc. **000**, 1–5 (2012) Printed 28 August 2012 (MN LaTeX style file v2.2)

arXiv:1208.5455v1 [astro-ph.HE] 27 Aug 2012

A determination of the masses of the components of Cyg X-3

Andrzej A. Zdziarski,[1] Joanna Mikołajewska[1] and Krzysztof Belczyński[2,3]
[1]*Centrum Astronomiczne im. M. Kopernika, Bartycka 18, PL-00-716 Warszawa, Poland*
[2]*Astronomical Observatory, University of Warsaw, Al. Ujazdowskie 4, 00-478 Warsaw, Poland*
[3]*Center for Gravitational Wave Astronomy, University of Texas at Brownsville, Brownsville, TX 78520, USA*

Submitted 2012 August 27

ABSTRACT
The Galactic X-ray binary Cyg X-3 consists of a compact object wind-fed by a Wolf-Rayet (WR) star. We determine its binary parameters and the mass-loss rate. Using the published infrared and X-ray data, we calculate the mass ratio and inclination. Then we fit the dependence of the mass-loss rate on the mass for WR stars of the WN type, and take into account a relation between the mass-loss rate and the binary period derivative, observed to be > 0 in Cyg X-3. This allows us to solve for the masses, inclination and mass-loss rate in the system. Our obtained mass-loss rate is almost identical to that from two independent estimates and consistent with other ones, which strongly supports the validity of this solution. We obtain the WR and compact object masses as $10.3^{+3.9}_{-2.8}{\rm M}_{\odot}$, $2.4^{+2.1}_{-1.1}{\rm M}_{\odot}$, respectively. Thus, either a neutron star or a low-mass black hole can be present. However, the radio, infrared and X-ray properties of the system suggest that the compact object is a black hole.

Key words: binaries: close – stars: individual: Cyg X-3 – stars: winds, outflows – stars: Wolf-Rayet – X-rays: binaries.

1 INTRODUCTION

In spite of its discovery already in 1966 (Giacconi et al. 1967), Cyg X-3 remains poorly understood. It is located at a distance of $d \simeq$ 7–9 kpc in the Galactic plane (Dickey 1983; Predehl et al. 2000; Ling, Zhang & Tang 2009). The donor appears to be a Wolf-Rayet (WR) star of the WN type (van Kerkwijk et al. 1992, 1996; van Kerkwijk 1993; Fender, Hanson & Pooley 1999). However, the compact object may be either a neutron star (NS) or a black hole (BH, e.g., Lommen et al. 2005; hereafter L05; Vilhu et al. 2009, hereafter V09).

Here, we self-consistently determine the binary parameters and the mass-loss rate, $\dot{M}$, of the donor. We use published results for the radial velocities and a relationship between the period derivative, $\dot{P}$, and the total mass of the system. We close the equations by finding a tight correlation between $\dot{M}$ and the WR mass, $M_{\rm WR}$.

2 THE BINARY PARAMETERS AND MASS-LOSS RATE

The best existing determination of the mass function of the compact object appears to be that of Hanson, Still & Fender (2000), who measured the radial velocity of the WR star as $K_{\rm WR} \simeq 109 \pm 13$ km/s. Then, V09 studied orbital modulation of X-ray lines in the system using *Chandra*, in particular those originating close to the compact object. In their determination of the radial velocity of the compact object, $K_{\rm C}$, they calculated the average based on 4, 8 and 10 bins. We note that in the case of 8 bins, the two with the maximum blueshift were removed. That case gave the radial velocity substantially less than the other ones, which reduced the resulting velocity average. Here, we use their values for the 4 and 10 bins with, in addition, the case of the *Chandra* observation of Obsid 6601 from V09. The unweighted average of the three measurements gives then the compact-object velocity of $K_{\rm C} \simeq 469 \pm 88$ km/s. We have also repeated their fits using their data from fig. 7, obtaining values very similar to their, and consistent with the above average. Thus, we use this value of $K_{\rm C}$ below.

With the binary period of $P = 0.19969$ d (Singh et al. 2002), the mass functions are $f_{\rm WR} = 2.13^{+1.44}_{-0.99}{\rm M}_{\odot}$ and $f_{\rm C} = 0.027^{+0.011}_{-0.008}{\rm M}_{\odot}$. (The original value of $K_{\rm C} \simeq 418 \pm 123$ km/s of V09 implies $f_{\rm WR} = 1.51^{+1.59}_{-0.92}{\rm M}_{\odot}$.) This (for our value of $f_{\rm WR}$) implies the mass ratio and the inclination of

$$q \equiv \frac{M_{\rm C}}{M_{\rm WR}} = \frac{K_{\rm WR}}{K_{\rm C}} \simeq 0.23^{+0.09}_{-0.06}, \qquad (1)$$

$$\sin i = \left[\frac{(K_{\rm WR}+K_{\rm C})^2 K_{\rm C} P}{2\pi G M_{\rm WR}}\right]^{1/3} \simeq (0.69 \pm 0.12)\left(\frac{M_{\rm WR}}{10{\rm M}_{\odot}}\right)^{-1/3}, \qquad (2)$$

respectively (note some misprints in the corresponding equations of V09). Here, we have estimated the errors as corresponding to the extrema of the velocities. At $M_{\rm WR} = 10{\rm M}_{\odot}$, $i \simeq 43^{\circ}{}^{+11}_{-9}$. If $i \geq 30^{\circ}$, which is likely given the strong IR and X-ray orbital modulation, the above constraints imply $M_{\rm WR} \lesssim 42{\rm M}_{\odot}$ and the compact-object mass of $M_{\rm C} \lesssim 9{\rm M}_{\odot}$.

Based on their population synthesis and evolution calculations, L05 claimed that the case of an NS can correspond only to the donor being a low-mass, $\lesssim 1.5{\rm M}_{\odot}$, He star accreting at a high rate onto the NS through a Roche-lobe overflow. Such an He star has a negligible wind and the observed WR features originate then in an outflow centered on the NS. Then, both IR and X-ray spectral

© 2012 RAS

features originate in the neutron-star surroundings. It is then not clear how to explain the orbital modulation of the continuum, the emission lines fluxes and the radial velocities. Specifically, both the lines originating in the WR-like outflow and around the NS should have similar dependencies on the orbital phase, contrary to the observations (Hanson et al. 2000; V09). On the other hand, V09 self-consistently explain the radial velocity and flux modulations with the orbital phase for both the X-ray and near-IR features (lines and continuum) in the frame of a model consisting of a real WR star and a compact companion. In particular, if the minimum flux (X-ray, near-IR) occurs when the WR transits in front of the compact object, the radial velocity curves tracing the WR star (He i absorption; Hanson et al. 2000) and the compact object (Fe xxvi; V09) have the minima/maxima at the respective quadratures, i.e., are shifted by 0.25*P* and 0.75*P*, respectively, whereas the radial velocities of the IR He ii emission formed in the X-ray shadow cone behind the WR star, where the wind is photo-ionized by its EUV radiation, show a direct correlation with the light curve. Such a behavior appears impossible to explain in terms of the model in which all WR features arise in the matter re-ejected from the region centered at the compact object. We note that this model was proposed before the results of V09 were obtained. Ruling it out, the mass loss is via the standard WR stellar wind.

We have then two additional constraints. The binary is detached (van Kerkwijk et al. 1992), and thus there is no Roche lobe overflow and accretion is from the wind. Most of the wind leaves the system carrying away its angular momentum. This leads to slowing down the binary rotation at a rate (e.g., Davidsen & Ostriker 1974),

$$\frac{\dot P}{P} \simeq \frac{2\dot M}{M}, \tag{3}$$

where $M = M_{\rm WR} + M_{\rm C}$. Ogley, Bell Burnell & Fender (2001) critically considered assumptions made in obtaining this relationship. Based on their discussion, there appear to be no major problems with this method when applied to Cyg X-3. The most recent published estimate for $\dot P/P$ is $(1.05 \pm 0.04) \times 10^{-6}$ yr^{-1} (Singh et al. 2002). It has been confirmed and improved by S. Kitamoto (private comm.) by including *Suzaku* observations, which yields $\dot P/P \simeq (1.03 \pm 0.02) \times 10^{-6}$ yr^{-1}. Then, $M \simeq 2\dot M P/\dot P \simeq 19 {\rm M}_\odot (\dot M/10^{-5} {\rm M}_\odot/{\rm yr})$.

The second constraint is of the mass-loss rate vs. mass relationship in WR stars. van Kerkwijk et al. (1996) classified Cyg X-3 as the WN 4–8 stellar type, which is intermediate between the WNE type, almost H free, and WNL, with substantial fraction of H, on the basis of the I and K-band emission lines. Thus, we consider here WN stars regardless of their H content, though the $\dot M(M_{\rm WR})$ relationship we find is similar if we restrict our sample to the He fraction of $Y \geq 0.92$. We use the sample of table 5 of Nugis & Lamers (2000, hereafter NL00), containing 34 WN stars (31 of them with the type WN 4–8), and giving $\dot M$ corrected for wind clumping (unlike a number of previous studies), which we show in Fig. 1. The metallicity of all these stars is close to solar, $Z = 0.0172$–0.0176. In the sample, 29 stars have $M_{\rm WR} < 22{\rm M}_\odot$, and 5 stars have $M_{\rm WR} = (40$–$57){\rm M}_\odot$. We can see that the $\dot M(M_{\rm WR})$ relationship is rather steep for the former but it saturates for the latter. Since our considerations above point to a relatively moderate mass of the WR star, we fit only the former subsample. Using a fitting method symmetric in $M_{\rm WR}$ and $\dot M$ (see appendix C of Zdziarski et al. 2011), we obtain

$$\dot M \simeq \dot M_0 \left(\frac{M_{\rm WR}}{\langle M\rangle}\right)^n, \quad \dot M_0 = (1.9 \pm 0.2) \times 10^{-5} \frac{{\rm M}_\odot}{\rm yr},$$

$$\langle M\rangle \simeq 14.7{\rm M}_\odot, \quad n \simeq 2.93 \pm 0.38, \quad M \lesssim 22{\rm M}_\odot, \tag{4}$$

shown in Fig. 1. Here, the normalization is at the average of the fitted $M_{\rm WR}$, $\langle M\rangle$. Equation (4) for a fiducial $\dot M = 10^{-5}{\rm M}_\odot/{\rm yr}$ implies $M_{\rm WR} \simeq (12 \pm 1){\rm M}_\odot$. NL00 fitted the entire WN sample, including the massive stars, and thus obtained a much flatter dependence, see their equation (23). However, it is clear from Fig. 1 that this is a very inaccurate description of the actual dependence. Note that clumping reduces $\dot M$ for a given $M_{\rm WR}$, and the above relationship is a factor of 2–3 below that of Langer (1989) or L05. On the other hand, our fit predicts $\dot M$ *higher* by a factor of 3.6 than that given by Vink & de Koter (2005) for the WN 8 star WR 40 (also present in the sample used here) with $M_{\rm WR} \simeq 20{\rm M}_\odot$, used as the normalization of their theoretical dependence of $\dot M(Z)$ for WN stars. This, however, appears to be due to a mistake in their coping that value of $\dot M$ from Herald, Hillier & Schulte-Ladbeck (2001), whose value of $3.1 \times 10^{-5}{\rm M}_\odot/{\rm yr}$ is almost the same as that of NL00, and consistent with our fit. We also note that although Herald et al. (2001) claim that their value of $\dot M$ for another WN 8 star, WR 16, is different from that of NL00 due to a different treatment of clumping, the difference is entirely accounted for by the different values of the used distance. Summarizing, we have not found any valid objections to the results of NL00 for their WN sample.



Figure 1. The mass-loss rates corrected for wind clumping vs. the mass for WN stars in the sample of NL00. The solid line shows the fit to all points with $M_{\rm WR} < 22{\rm M}_\odot$, and the dotted lines delineate the rms of these points. The red and blue crosses show the objects with reliable determinations of $M_{\rm WR}$ and $\dot M$ and those with only rough estimates (according to NL00), respectively. We see no systematic difference between these two categories at $M_{\rm WR} < 22{\rm M}_\odot$.

Equations (3) and (4) can be solved for $M_{\rm WR}$ and $\dot M$ as functions of q, equation (1),

$$M_{\rm WR} = \left[\frac{\dot{P}(1+q)\langle M\rangle^n}{2P\dot{M}_0}\right]^{\frac{1}{n-1}}, \tag{5}$$

$$\dot{M} = \dot{M}_0^{-\frac{1}{n-1}}\left[\frac{\dot{P}(1+q)\langle M\rangle}{2P}\right]^{\frac{n}{n-1}}. \tag{6}$$

Using the $\dot{P}/P$ of Kitamoto, these formulae give $M_{\rm WR} \simeq 10.3^{+1.5}_{-1.9}{\rm M}_\odot$, $M_{\rm C} \simeq 2.4^{+1.4}_{-1.0}{\rm M}_\odot$, $\dot{M} \simeq 6.5^{+1.7}_{-1.5} \times 10^{-6}{\rm M}_\odot$/yr, and $i = 43^{\circ}{}^{+14}_{-9}$ [from equation (2)]. The lower limits on the masses and $\dot{M}$ and the upper limit on i correspond to the lower limits on q, $\dot{P}$ and n and the upper limit on $\dot{M}_0$, and vice versa. Using the original mass function of V09 changes these results only slightly. We note here that the fitting method finds the uncertainty of the *average* dependence. Its spread, or rms, is much larger, and it is partly intrinsic and partly due to measurement errors. Without detailed knowledge of the errors, we can make a conservative assumption that the observed dispersion of $\dot{M}$ for a given $M_{\rm WR}$ is mostly intrinsic, and due to the internal structure and/or Y affecting $\dot{M}$. An estimate of the rms is $\Delta\dot{M}_0\sqrt{N-1}$, where $N = 29$ is the number of stars in the fitted sample. We then keep n fixed, which corresponds to moving up and down the best fit dependence within the observed spread, as illustrated by the dotted lines in Fig. 1. This way of estimating the $\dot{M}(M_{\rm WR})$ uncertainty yields somewhat wider confidence limits of $M_{\rm WR} \simeq 10.3^{+3.9}_{-2.8}{\rm M}_\odot$, $M_{\rm C} \simeq 2.4^{+2.1}_{-1.1}{\rm M}_\odot$, $i = 43^{\circ}{}^{+17}_{-12}$, $\dot{M} \simeq 6.5^{+3.3}_{-2.1}\times 10^{-6}{\rm M}_\odot$/yr. Concluding, both ways of estimating the uncertainties allow either a low-mass BH or an NS to be present.

We note that Singh et al. (2002) and Kitamoto (priv. comm.) also fitted cubic ephemerides to their data, i.e., allowed for $\ddot{P} \neq 0$. They obtained rather small values of $\dot{P}/\ddot{P} \simeq -60 \pm 70$ yr and -130 ± 80 yr, respectively. These ephemerides would thus predict that $\dot{P}$ changes its sign in $\sim 10^2$ yr, which is obviously $\ll$ any evolutionary time scale of the wind. Thus, this model is inconsistent with the $\dot{P}$ being due to the loss of the angular momentum via wind, as assumed in equation (3). If such an ephemeris were shown to be correct, another explanation, e.g., a third body, would be required. However, a linear + sinusoidal ephemeris gives a very bad fit to those data. Moreover, a cubic ephemeris for the most recent data set yields no improvement of the fit over the parabolic one, and the same reduced χ^2 (Kitamoto, priv. comm.). We thus do not consider this ephemeris any more.

We then compare our results on $\dot{M}$ with other independent estimates. The most detailed study of the constraint from X-ray absorption appears to be that of Szostek & Zdziarski (2008). They took into account wind clumping as well as the presence of a homogeneous wind phase. They obtained results for the cases of a low and high mass of the WR star, $6.4{\rm M}_\odot$ and $70{\rm M}_\odot$, respectively. Our range of $M_{\rm WR}$ is much closer to the lower one, and the dependence on $M_{\rm WR}$ is relatively modest. For the low-mass case, they obtained $\dot{M} \simeq (6\text{–}8) \times 10^{-6}{\rm M}_\odot$/yr, almost the same as our values obtained here. A related (but independent) estimate is that of Zdziarski et al. (2012), who studied X-ray orbital modulation in Cyg X-3. Its orbital-phase dependence in hard X-rays, where the opacity is likely to be dominated by Thomson scattering, implies $\dot{M} \simeq (7\text{–}8) \times 10^{-6}{\rm M}_\odot$/yr for the terminal wind velocity of 1500–1700 km/s (Fender et al. 1999; van Kerkwijk et al. 1996). Then, Waltman et al. (1996) obtained $\dot{M} \lesssim 1.0\times 10^{-5}{\rm M}_\odot$/yr from the radio delays interpreted as being due to wind opacity. This is again fully consistent with our values. The above methods are independent of the distance to Cyg X-3.

On the other hand, van Kerkwijk (1993) and Ogley et al. (2001) obtained higher values of $\dot{M}$ by interpreting the observed IR fluxes as solely due to wind emission. van Kerkwijk (1993) obtained $\dot{M} \simeq 4 \times 10^{-5}{\rm M}_\odot$/yr without taking into account clumping of the wind. Including it lowers $\dot{M}$ by a factor of several (Szostek & Zdziarski 2008) and thus makes it consistent with our present results. Ogley et al. (2001) obtained $\dot{M} \simeq (4\text{–}30) \times 10^{-5}{\rm M}_\odot$/yr using different assumptions about the wind, but also neglecting clumping. The lowest value is for a strongly ionized wind. Using that assumption and taking into account clumping can again reconcile their results with ours. Furthermore, the wind-emission calculations depend on the distance as $\dot{M} \propto d^{3/2}$, and the above authors assumed $d \simeq 10$ kpc. If $d \simeq 7$ kpc, their values of $\dot{M}$ should be reduced by ∼40 per cent. In addition, there can be a substantial IR contribution from the synchrotron emission of the jets, which would again reduce the calculated $\dot{M}$. Summarizing, all the published estimates of $\dot{M}$ are compatible with our present results.

3 DISCUSSION

As we noted above, L05 claimed that only a BH can be present in the case of a WR donor, based on their population synthesis results. However, given uncertainties intrinsic to the population-synthesis method, this conclusion appears uncertain. Also, their specific version of this method suffers from a number of problems. One of them is their condition for accretion disc formation for an isotropic wind of Ergma & Yungelson (1998). However, if the wind velocity at the Roche-lobe surface of the WR donor is less than the escape velocity from that surface, the wind will be strongly focused towards the accretor, which facilitates accretion disc formation. Numerical simulations for such a wind Roche-lobe overflow (Mohamed & Podsiadlowski 2012) indicate that this mass transfer mechanism may occur in systems where the wind acceleration zone lies close to or makes a significant fraction of the Roche-lobe radius. This is very likely to be the case in Cyg X-3 given its short orbital period. Thus, our conclusion above that an NS *may* be present in Cyg X-3 appears to be valid.

On the other hand, there is a lot of other (though still circumstantial) evidence for the compact object in Cyg X-3 being a BH. Its spectral states have spectra showing a general resemblance to those of the canonical spectral states of BH binaries (Szostek & Zdziarski 2008; Zdziarski, Misra & Gierliński 2010). The existing differences may be explained by scattering in the very strong stellar wind in Cyg X-3 (Zdziarski et al. 2010). The radio flux in the hard spectral state is strongly correlated with that in soft X-rays, which correlation is very similar (Gallo, Fender & Pooley 2003; Szostek, Zdziarski & McCollough 2008) to that seen in BH X-ray binaries. The normalization of the radio flux in Cyg X-3 is somewhat above that for BH binaries, which excess can be due to strong absorption of soft X-rays in Cyg X-3. On the other hand, NS X-ray binaries in the analogous state have the radio flux $\gtrsim 30$ times weaker on average (Migliari & Fender 2006), in stark contrast to the case of Cyg X-3. As pointed out by Szostek & Zdziarski (2008), all NSs in confirmed high mass X-ray binaries are highly magnetized (which is related to them being young systems), and are usually X-ray pulsars. Even in cases with no detected X-ray pulsations, their bremsstrahlung type spectra bear no resemblance to Cyg X-3, which is also a young system. Then, V09 fitted the IR-flux orbital modulation and found it is compatible with a low inclination, $i \sim 30^\circ$, but not with $i \sim 60^\circ$, implying the compact object to be a BH. A low inclination, $i \simeq 30\text{–}40^\circ$, has also been found by Zdziarski et al. (2012) as accounting for the orbital modulation of the X-ray continuum.

Following the above circumstantial evidence and our estimate of the compact object mass ($M_{\rm C} = 2.4^{+2.1}_{-1.1}{\rm M}_{\odot}$) we argue that Cyg X-3 hosts a low mass BH. The actual BH mass is most likely in the range ($M_{\rm NS,max}$–$4.5{\rm M}_{\odot}$), where $M_{\rm NS,max}$ is the still unknown maximum NS mass. The recent precise mass measurements of Galactic pulsars require that $M_{\rm NS,max} > 2{\rm M}_{\odot}$ (Demorest et al. 2010), while dynamical BH mass estimates impose that $M_{\rm NS,max} < 5{\rm M}_{\odot}$ (Özel et al. 2010; Farr et al. 2011). Theoretical calculations of dense nuclear matter indicate that $M_{\rm NS,max} < 2.4{\rm M}_{\odot}$ (Lattimer & Prakash 2010). When we impose above limits on our estimate we obtain $M_{\rm BH} = 2.4^{+2.1}_{-0.4}{\rm M}_{\odot}$. This puts the BH in Cyg X-3 right into the mass gap, the dearth of compact objects in the (2–5)${\rm M}_{\odot}$ mass range, first noted by Bailyn et al. (1998).

The existence of the mass gap was recently discussed in context of supernova explosion models (Belczyński et al. 2012). For the rapidly developing explosions (within ∼200 ms after the core bounce), the mass gap appears in simulations as observed for the Galactic BH X-ray binaries. However, there is a small addition of a few systems with compact objects in (2–3.5)${\rm M}_{\odot}$ mass range (see fig. 1 of Belczyński et al. 2012). These have formed via accretion onto an NS (originally formed in the supernova explosion) from a binary companion. These compact objects are most likely light BHs formed via accretion induced collapse of the NS (van den Heuvel & De Loore 1973).

The mass gap was also recently interpreted as a potential observational artefact caused by the systematic uncertainties in dynamical BH mass measurements (Kreidberg et al. 2012). No mass gap scenario corresponds to delayed supernova models (developing on a timescale ∼500–1000 ms) that produce compact objects within (2–5)${\rm M}_{\odot}$ mass range (Belczyński et al. 2012). In this case, the light BH in Cyg X-3 most likely have formed in supernova explosion from an $M_{\rm zams} = (20$–$40){\rm M}_{\odot}$ progenitor star (Fryer et al. 2012). As long as there is no consensus on the mass gap existence, both formation scenarios, either via accretion induced collapse or in the delayed supernova explosion, are to be considered.

The Eddington luminosity for He accretion is $L_{\rm E} \simeq 2.5 \times 10^{38}(M_{\rm C}/{\rm M}_{\odot})$ erg s^{-1}. The absorption corrected, phase-averaged, bolometric fluxes in the hard and soft spectral state (assuming the low-mass model of Szostek & Zdziarski 2008; the hard-state flux corrected in Szostek et al. 2008) are 8.5×10^{-9} erg cm^{-2} s^{-1} and 2.7×10^{-8} erg cm^{-2} s^{-1}, respectively. These are orbital-phase averages; they should be increased by a factor $\simeq 1.5$ and 2 in the hard and soft state, respectively, to obtain the intrinsic values (Zdziarski et al. 2012). Assuming $d = 7$ kpc, we obtain the bolometric hard and soft state luminosities of $\simeq 7 \times 10^{37}$ erg s^{-1} and $\simeq 3 \times 10^{38}$ erg s^{-1}, respectively. At $M = 4{\rm M}_{\odot}$, they correspond to $\simeq 0.07L_{\rm E}$ and $\simeq 0.3L_{\rm E}$, respectively. These Eddington ratios are within the ranges observed for black-hole binaries in these two states (Done, Gierliński & Kubota 2007).

4 CONCLUSIONS

We have have determined the masses of the components, the inclination and the donor mass-loss rate in the Cyg X-3 binary. We have used the $K_{\rm C}$ measurement of Hanson et al. (2000) and revised the determination of $K_{\rm WR}$ of V09 based on their data. We then assumed that $\dot{M}$ is related to $\dot{P}$ and the total mass via the orbital angular momentum being lost by the wind. We have also substantially revised existing dependencies of $\dot{M}(M_{\rm WR})$ for WN stars by using the clumping-corrected results of NL00 and fitting them for moderate masses only. The resulting four equations are solved for $M_{\rm WR}$, $M_{\rm C}$, i and $\dot{M}$. The obtained $\dot{M}$ is in a very good agreement with two independent calculations, and are consistent with all other determinations, provided those based on the IR fluxes are corrected for clumping.

Our results for the (conservative) assumption that the observed spread of $\dot{M}(M_{\rm WR})$ is intrinsic are $M_{\rm WR} \simeq 10.3^{+3.9}_{-2.8}{\rm M}_{\odot}$, $M_{\rm C} \simeq 2.4^{+2.1}_{-1.1}{\rm M}_{\odot}$, $i = 43^{\circ}{}^{+17}_{-12}$, $\dot{M} \simeq 6.5^{+3.3}_{-2.1} \times 10^{-6}{\rm M}_{\odot}$/yr. We argue that although an NS is allowed to be present, a BH is more likely given the radio, IR and X-ray properties of Cyg X-3.

ACKNOWLEDGMENTS

This research has been supported in part by the Polish NCN grants N N203 581240, N N203 404939, 362/1/N-INTEGRAL/2008/09/0, and the NASA Grant NNX09AV06A to the UTB Center for Gravitational Wave Astronomy (KB). We thank Janusz Ziółkowski for a valuable discussion.

REFERENCES

Bailyn C. D., Jain R. K., Coppi P., Orosz J. A., 1998, ApJ, 499, 367

Belczyński K., Wiktorowicz G., Fryer C., Holz D., Kalogera V., 2012, ApJ, in press, arXiv:1110.1635

Davidsen A., Ostriker J. P., 1974, ApJ, 189, 331

Demorest P. B., Pennucci T., Ransom S. M., Roberts M. S. E., Hessels J. W. T., 2010, Natur, 467, 1081

Dickey J. M., 1983, ApJ, 273, L71

Done C., Gierliński M., Kubota A., 2007, A&ARv, 15, 1

Ergma E., Yungelson L. R., 1998, A&A, 333, 151

Farr W. M., Sravan N., Cantrell A., Kreidberg L., Bailyn C. D., Mandel I., Kalogera V., 2011, ApJ, 741, 103

Fender R. P., Hanson M. M., Pooley G. G., 1999, MNRAS, 308, 473

Fryer C. L., Belczyński K., Wiktorowicz G., Dominik M., Kalogera V., Holz D. E., 2012, ApJ, 749, 91

Gallo E., Fender R. P., Pooley G. G., 2003, MNRAS, 344, 60

Giacconi R., Gorenstein P., Gursky H., Waters J. R., 1967, ApJ, 148, L119

Hanson M. M., Still M. D., Fender R. P., 2000, ApJ, 541, 308

Herald J. E., Hillier D. J., Schulte-Ladbeck R. E., 2001, ApJ, 548, 932

Kreidberg L., Bailyn C., Farr W., Kalogera V. 2012, ApJ, submitted, arXiv:1205.1805

Langer N., 1989, A&A, 220, 135

Lattimer J. M., Prakash M., 2010, In Lee S., ed., Gerry Brown's Festschrift. World Scientific, arXiv:1012.3208

Ling Z., Zhang S. & Tang S., 2009, ApJ, 695, 1111

Lommen D., Yungelson L., van den Heuvel E., Nelemans G., Portegies Zwart S., 2005, A&A, 443, 231 (L05)

Migliari S., Fender R. P., 2006, MNRAS, 366, 79

Mohamed S., Podsiadlowski P., 2012, BaltA, 21, 88

Nugis T., Lamers H. J. G. L. M., 2000, A&A, 360, 227 (NL00)

Ogley R. N., Bell Burnell S. J., Fender R. P., 2001, MNRAS, 322, 177

Özel F., Psaltis D., Narayan R., McClintock J. E., 2010, ApJ, 725, 1918

Predehl P., Burwitz V., Paerels F., Trümper J., 2000, A&A, 357, L25

Schaerer D., Maeder A., 1992, A&A, 263, 129

Singh N. S., Naik S., Paul B., Agrawal P. C., Rao A. R., Singh K. Y., 2002, A&A, 392, 161
Szostek A., Zdziarski A. A., 2008, MNRAS, 386, 593 (SZ08)
Szostek A., Zdziarski A. A., McCollough M. L., 2008, MNRAS, 388, 1001
van den Heuvel E. P. J., De Loore C., 1973, A&A, 25, 387
van Kerkwijk M. H., 1993, A&A, 276, L9
van Kerkwijk M. H., et al., 1992, Natur, 355, 703
van Kerkwijk M. H., Geballe T. R., King D. L., van der Klis M., van Paradijs J., 1996, A&A, 314, 521
Vilhu O., Hakala P., Hannikainen D. C., McCollough M., Koljonen K., 2009, A&A, 501, 679 (V09)
Vink J., de Koter A., 2005, A&A, 442, 587
Waltman E. B., Foster R. S., Pooley G. G., Fender R. P., Ghigo F. D., 1996, AJ, 112, 2690
Zdziarski A. A., Misra R., Gierliński M., 2010, MNRAS, 402, 767
Zdziarski A. A., Skinner G. K., Pooley G. G., Lubiński P., 2011, MNRAS, 416, 1324
Zdziarski A. A., Maitra C., Frankowski A., Skinner G. K., Misra R., 2012, MNRAS, in press, arXiv:1205.4402